

# UNCLASSIFIED

AD NUMBER	
AD334076	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	confidential
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 24 SEP 1946. Other requests shall be referred to Naval Ordnance Lab., White Oak, MD.
AUTHORITY	
30 Sep 1974, DoDD 5200.10; USNSWC ltr, 7 Oct 1974	

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD. 334076

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA

CLASSIFICATION CHANGED  
TO UNCLASSIFIED  
FROM CONFIDENTIAL  
PER AUTHORITY LISTED IN

Bul. no. 68-11

1 June 1968



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

NOLTR 62-175

CATALOGED BY ASTIA

AS AD NO. ~~334 076~~


334 076

HEAT RESISTANT EXPLOSIVES. XV. (U)

Alternate Synthetic Routes to 3,3'-Diamino-  
2,2',4,4',6,6'-hexanitrobiphenyl, DIPAM (C)

NOL

24 SEPTEMBER 1962



UNITED STATES NAVAL ORDNANCE LABORATORY, WHITE OAK, MARYLAND

NOLTR 62-175

NOTICE: This material contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C. Sections 793 and 794, the transmission or revelation of which in any manner to an unauthorized person is prohibited by law.

Downgraded at 3 Year Intervals  
Declassified after 12 Years. DOD Dir 5200.10

Best Available Copy

~~CONFIDENTIAL~~  
NOLTR 62-175

HEAT RESISTANT EXPLOSIVES. XV. (U)  
Alternate Synthetic Routes to 3,3'-  
Diamino-2,2',4,4',6,6'-hexanitrobiphenyl,  
DIPAM (C)

Prepared by: Robert E. Oesterling  
Joseph C. Dacons  
Lloyd A. Kaplan

ABSTRACT: DIPAM has been prepared from m-bromoanisole by a new three step synthesis involving the following reactions: m-bromoanisole to trinitro-m-bromoanisole which was in turn converted to dimethoxy hexanitrobiphenyl via the Ullmann reaction. Conversion of the dimethoxy derivative to DIPAM was readily achieved by treating it with ammonia in alcohol.

A second route, starting with m-bromoaniline via the sequence methyl m-bromophenylurethane to methyl trinitro-m-bromophenylurethane to the bis-methyl urethane of DIPAM to DIPAM was also attempted because of its simplicity. However the sole product isolated from this sequence was picramide. (C)

Approved by:

Darrell V. Sickman, Chief  
Organic Chemistry Division  
Chemistry Research Department  
U. S. NAVAL ORDNANCE LABORATORY  
White Oak, Silver Spring, Maryland

~~CONFIDENTIAL~~

NOLTR 62-175

24 September 1962

This report describes a new method for the preparation of DIPAM, a new thermally stable high explosive which may be useful in mild detonating fuse for stage separation in advanced POLARIS missiles. This compound was previously described in NOLTR 62-82 which was Part XIII in the series Heat Resistant Explosives. The work described herein was carried out under Task FR-44.

R. E. ODENING  
Captain, USN  
Commander

*Albert Lightbody*  
ALBERT LIGHTBODY  
By direction

~~CONFIDENTIAL~~

CONFIDENTIAL  
NOLTR 62-175

TABLE OF CONTENTS

	Page
INTRODUCTION .....	1
DISCUSSION	
m-Bromoaniline to DIPAM .....	3
m-Bromoanisole to DIPAM .....	5
Resorcinol Monomethyl Ether to DIPAM .....	
EXPERIMENTAL .....	7
REFERENCES .....	11

# HEAT RESISTANT EXPLOSIVES. XV.

## INTRODUCTION

In a previous NOLTR (1) the preparation of a new, thermally stable high explosive 3,3'-diamino-2,2',4,4',6,6'-hexanitro-biphenyl, DIPAM, was described. This explosive was particularly interesting since not only was it relatively non-volatile at elevated temperatures and reduced pressures, but it was also found that its detonation would propagate in very small diameters and around corners. These latter two properties led to the testing of DIPAM for application in mild detonating fuse for use in stage separation in advanced Polaris missiles.

The preparation of DIPAM described previously (1) involved a six step synthesis which is described schematically in Fig. 1.




Figure 1



CONFIDENTIAL  
NOLTR 62-175

m-Bromotoluene was nitrated in a one step nitration with ninety percent nitric acid and 30 percent oleum by heating at reflux for several hours after first maintaining the mixture at 55° for several hours. The product II, generally obtained in recrystallized yields of 86 percent, was isolated by drowning in ice.

The conversion of the trinitro derivative II to 3,3'-dimethyl-2,2',4,4',6,6'-hexanitrobiphenyl, III, was accomplished by the Ullmann method which consists of heating a halogenated benzene derivative at about reflux temperatures in nitrobenzene with "activated" copper powder. In this instance, the procedures of Moyer, Bock, and Adams (2) were followed. The biphenyl III was generally isolated in yields of 77 percent after crystallization.

Oxidation of the biphenyl III to dipicric acid, IV, was accomplished by the procedure of Moyer, Bock, and Adams (2) utilizing a mixture of sodium dichromate, nitric and sulfuric acids. This procedure was shown (3) to yield dipicric acid and not the reported (2) 3,3'-dicarboxy-2,2',4,4',6,6'-hexanitrobiphenyl. The yields of IV obtained by this procedure were generally about 75 percent.

Conversion of IV to the dichloro derivative V was accomplished by reacting the dipyridinium salt of IV with phosphorus oxychloride. This technique involved treating IV with pyridine to form the dipyridinium salt which was isolated as a crystalline material in yields of 89 percent. The dipyridinium salt was then added to an excess of phosphorus oxychloride, after which the mixture was drowned in ice. This procedure afforded the dichloro derivative V in average yields of 99 percent from the dipyridinium salt.

Finally, conversion of the dichloro derivative V to DIPAM, VI, was readily accomplished by adding the dichloro derivative to a solution of ammonia in ethanol at 10°. A 90 percent yield of DIPAM was obtained upon recrystallization from acetone-ethanol.

The above sequence gave an overall yield of DIPAM of 39.4 percent. Although this synthesis was attractive from the standpoint of the rather excellent yields obtained in most of the steps, the multiplicity of procedures involved in the sequence is not only somewhat time consuming but also tends to reduce the overall yield of DIPAM obtained. Thus, it appeared reasonable to look at some other possible synthetic routes to DIPAM which would involve fewer steps.

### DISCUSSION

Two possible alternate routes presented themselves for consideration. The first of these started with m-bromoaniline which was converted to the methyl urethane VII. Nitration of VII with mixed acid would yield the trinitro derivative VIII. Reacting VIII under Ullmann conditions, hopefully, would afford the bis-methyl urethane of DIPAM, IX, which could be readily hydrolyzed to DIPAM with sulfuric acid. This sequence is described in Figure 2.




Figure 2

The interesting features of the above sequence are that it starts with the readily available m-bromoaniline\* and all of the steps with the exception of the Ullmann reaction leading to the

\* m-Bromoaniline can be readily prepared from m-bromonitrobenzene; the corresponding chloro compounds are available in commercial quantities.

CONFIDENTIAL  
NOLTR 62-175

biphenyl derivative IX have been carried out previously (4). Its disadvantage lies in the fact that the Ullmann reaction, when carried out with compounds containing "active hydrogen atoms", viz. carboxylic acids, phenols, etc., is reported to yield reduction products (5). Thus the reaction of the trinitro derivative VIII under Ullmann conditions might yield the methyl urethane of picramide as the sole product. This would perhaps occur by reaction of the "organo-copper" intermediate with the acidic hydrogen atom of the amine function of a second molecule. Hydrolysis of this reduction product with sulfuric acid would in turn afford picramide instead of DIPAM.

Conversion of m-bromoaniline to the methyl urethane VII was accomplished in 82 percent yield by refluxing the amine and methyl chloroformate in benzene with sodium carbonate. Nitration of the methyl urethane VII to the trinitro derivative VIII was readily effected in 66 percent yield by a modification of the procedure of de Monchy (4). When the trinitro derivative was subjected to Ullmann conditions, it was observed that reaction set in at about 150° as evidenced by the disappearance of the bright copper color in the reaction mixture. The reaction proceeded smoothly at this temperature and the solid material at the end of the reaction was found to consist of unreacted copper and cuprous halide. The organic product from this reaction, obtained in low yield, could not be crystallized to yield a sharp melting product. Therefore it was hydrolyzed directly with sulfuric acid to remove the urethane function. Workup of the hydrolysis mixture yielded picramide and not DIPAM, thus indicating that the Ullmann reaction did not take the normal path, but instead effected reduction of the trinitro-m-bromophenylurethane, VIII.

The second route selected did not suffer from the possibility that the Ullmann reaction would yield reduction products. This sequence, Figure 3, starts with the conversion




Figure 3

of m-anisidine, X, to m-bromoanisole, XI\*, by means of the usual Sandmeyer reaction. In this instance, in a 75 percent yield. Nitration of XI to the trinitro derivative XII was effected in 74 percent yield (recrystallized, 88 percent crude) by a modification of the procedure of de Monchy (4). Reacting the trinitro derivative XII with activated copper powder under Ullmann conditions afforded a 78 percent recrystallized yield of the expected 3,3'-dimethoxy-2,2',4,4',6,6'-hexanitrobiphenyl, XIII. Finally, the dimethoxy derivative XIII was readily converted to DIPAM in 91 percent yield by treating it with alcoholic ammonia. The DIPAM so obtained melted at 295-96° without recrystallization; an authentic sample melted at 296-97° (simultaneous determination) and gave no depression on admixture with the product obtained by the above sequence of reactions.

\*m-Bromoanisole can also be prepared in 75 percent yield by the methylation of m-bromophenol (6). However, m-bromophenol is about four times more expensive than m-anisidine.

CONFIDENTIAL  
NOLTR 62-175

Thus, it was possible to prepare DIPAM by a four-step synthesis starting from m-anisidine\* in a 39.4 percent overall yield.

It should be noted that the above sequence leading to DIPAM was carried out in such a manner as to test the feasibility of the method. No attempt was made to optimize reaction conditions so as to obtain the maximum possible yields in each of the steps in the synthetic path. Undoubtedly, higher yields could be obtained in the nitration of m-bromoanisole to the trinitro derivative XII as evidenced by the 14 percent loss on purification of the nitration product. It is also probable that the yields obtained in the conversion of m-anisidine to m-bromoanisole and the coupling of the trinitro derivative XII to the biphenyl XIII could be increased somewhat by optimizing these reaction conditions.

A modification of the synthetic path described in Figure 3 starts with the more readily available resorcinol monomethyl ether. Nitration of the monomethyl ether of resorcinol under the appropriate conditions \*\* would afford 3-methoxy-2,4,6-trinitrophenol. Conversion of 3-methoxy-2,4,6-trinitrophenol to the chlorine analog of the trinitro derivative XII could probably be accomplished by means of the pyridine-phosphorus oxychloride method previously described (1). The chlorine analog of XII, 3-chloro-2,4,6-trinitroanisole, would then be reacted with copper powder under Ullmann conditions to yield the biphenyl XIII which in turn would be aminated to DIPAM with ammonia in alcohol.

Although this sequence requires the same number of steps as the route starting with m-anisidine, resorcinol monomethyl ether costs about one half as much as m-anisidine in research quantities\*\*\*. Therefore, the per pound cost of DIPAM

---

\* m-Anisidine is currently available in research quantities at a price of \$16.00 per 100 grams. It could undoubtedly be made more cheaply on a larger scale by the sequence: m-nitroaniline to m-nitrophenol to m-nitroanisole which can be reduced in 80 percent yield to m-anisidine (7).

\*\*The conditions used for this nitration would undoubtedly be similar to those used in the nitration of phenol to picric acid or resorcinol dimethyl ether to trinitroresorcinol dimethyl ether (8).

\*\*\*In large scale production, the cost of resorcinol monomethyl ether would be reduced even more than that of m-anisidine since its preparation involves a one-step reaction starting with \$0.75 a pound resorcinol.

CONFIDENTIAL  
NOLTR 62-175

prepared by this route would be less than that prepared by the sequence described in Figure 3. This proposed synthesis will be tested in the future and the results will be the subject of a forthcoming NOLTR.

EXPERIMENTAL

m-Bromoanisole (XI) was prepared by a modification of the method used for the preparation of p-bromotoluene from p-toluidine (9). In the preparation of XI an equivalent amount of hydrobromic acid was used in place of sulfuric acid for the diazotization of m-anisidine. From 0.32 mole of m-anisidine there was obtained 0.24 mole (45 g - 75%) of m-bromoanisole; b. 209-12°.

3-Bromo-2,4,6-trinitroanisole (XII) was prepared by adding 0.1 mole (18.7 g) of m-bromoanisole dropwise to a stirred mixture of forty grams of 100% nitric acid and 50 ml of 30% oleum at 35 to 45°. During the addition of the anisole, the yellow trinitro derivative XII separated as a solid from the nitrating medium. After the addition of the anisole was completed, the mixture was stirred for an additional 20 minutes at 30 to 40°.

The reaction mixture was then poured into 500 ml of ice and water and the yellow granular solid that separated was filtered off with suction after all of the ice had melted. The crude product was washed several times with water and then dried in air to give 28.5 g (88%) of crude XII. Recrystallization from hexane-benzene gave 23.5 g (74%) bright yellow-orange crystals; m. 96-7° (lit. (10) 97°).

Anal. calc'd for  $C_7H_4BrN_3O_7$ ; C, 26.08; H, 1.24; N, 13.04.

Found; C, 26.30; H, 1.34; N, 12.81  $\lambda_{max}^{MeOH}$  (ε), 247 (13,880).

3,3'-Dimethoxy-2,2',4,4',6,6'-hexanitrobiphenyl (XIII) was prepared by heating a solution of 0.05 mole (16.1 g) of 3-bromo-2,4,6-trinitroanisole in 50 ml of nitrobenzene to 155°. After this temperature had been reached, 7 g of activated copper powder (1) was added in small portions with continual stirring over a period of 30 minutes while maintaining the temperature at 160-65°. The dark brown reaction mixture was stirred for an additional 15 minutes at this temperature after the addition of the copper was completed.

At the end of this time, the mixture was filtered with suction while hot and the solids washed thoroughly with acetone. The combined filtrates and washings were steam distilled until nitrobenzene could no longer be detected in the distillate.

CONFIDENTIAL  
NOLTR 62-175

After decanting the water from the residue remaining in the flask, it was dissolved in 200 ml of acetone, treated with Darco charcoal, and evaporated to a volume of about 50 ml. Water was added to the hot concentrate till the cloud point and the solution cooled to 0°. The yellow solid that separated was filtered and dried in air. It was purified further by redissolving it in 50 ml of acetone, treating with Darco, and adding 50 ml of ethanol. The resulting solution was concentrated by boiling until precipitation begins. The mixture was then cooled to 0° and filtered to yield 9.5 g (78%) of 3,3'-dimethoxy-2,2',4,4',6,6'-hexanitrobiphenyl; m. 187-90°. Recrystallization from benzene gave pale yellow crystals; m. 194-95°. The product melting at 190° was considered sufficiently pure for the amination step which follows.

Anal. calc'd. for  $C_{14}H_6N_6O_4$ ; C, 34.71; H, 1.65; N, 17.4.

Found; C, 34.92; H, 2.98; N, 17.9.  $\lambda_{\text{max}}^{\text{MeOH}}$  (ε), 243 (28,900).

Amination of 3,3'-dimethoxy-2,2',4,4',6,6'-hexanitrobiphenyl (XIII) to DIPAM (VI). Dry, gaseous ammonia was passed into a stirred mixture of 0.00414 mole (2.00 g) of the dimethoxy derivative XIII in 25 ml of absolute ethanol at room temperature. The mixture became dark red at first and then an orange product separated out. When an excess of ammonia was apparently present, the ammonia flow was stopped and the mixture boiled for ten minutes and then cooled to 0° in ice. The material which separated was filtered off with suction to yield 1.7 g (91%) of a yellow granular solid. After washing with ethanol, it was dried in vacuo at 100°; m. 295-96°. Authentic sample of DIPAM; m. 296-97° (simultaneous determination). The DIPAM obtained in this experiment gave no depression in melting point when mixed with an authentic sample.

Methyl 3-bromophenylurethane (VII) was prepared by refluxing a mixture of 0.22 mole (20.8 g) of methyl chloroformate, 0.20 mole (34.4 g) of m-bromoaniline, and 0.4 mole (42.2 g) of anhydrous sodium carbonate in 500 ml of benzene for 2.5 hours. The mixture was initially quite thick (amine hydrochloride) and as the reaction proceeded, thinned out to a suspension of inorganic salts in benzene. After the requisite time under reflux, the mixture was filtered hot and the filtrates were evaporated to dryness in vacuo. The residue was recrystallized from methanol-hexane to yield 37.7 g (82%) of white plates melting at 86-6.5° (lit. (4) 84°).

CONFIDENTIAL  
NOLTR 62-175

Methyl 3-bromo-2,4,6-trinitrophenylurethane (VIII) was prepared by a modification of the procedure of de Monchy (4). One tenth of a mole (23 g) of methyl 3-bromophenylurethane was dissolved in 50 ml of concentrated sulfuric acid while maintaining the temperature at 10° or lower. There was apparently no hydrolysis of the urethane function during this step as evidenced by the lack of carbon dioxide evolution from the solution. The resulting solution was added to a previously prepared mixture of 138 ml of concentrated sulfuric acid and 138 ml of 100% nitric acid which had been cooled to 0-10° in ice. The rate of addition of the urethane solution was such that the temperature was maintained between 5 and 10° during the addition. Following the completion of the addition, the resulting orange solution was stirred at 15-20° for 0.75 hour. During this period, a white solid separated (trinitro derivative VIII) and the mixture turned progressively yellower.

At the end of this time, the mixture was poured onto an excess of ice with stirring, and after the ice had melted, the solid which had separated was collected on a sintered glass funnel. The pale yellow solid was washed with water until neutral and recrystallized from methanol. There was obtained 24.1 g (66%) of an almost white solid; m. 226-228° uncorr. (lit. (4) 230°).

Attempted preparation of 2,2',4,4',6,6'-hexanitro-3,3'-di(carbamylmethyl)biphenyl (IX). In a one-liter flask fitted with a thermometer, stirrer, and solids addition funnel was placed 0.04 mole (14.6 g) of methyl 3-bromo-2,4,6-trinitrophenylurethane and 100 ml of nitrobenzene. The mixture was heated to 120° and a water pump vacuum was applied to distill off part (5 to 10 ml) of the nitrobenzene. This operation was carried out to "dry" the reaction mixture.

The solids addition funnel was placed back in the flask and a one gram portion of activated copper (1) from the total amount, 0.16 mole (10.16 g), was added. The temperature of the reaction mixture was raised slowly until evidence that reaction was occurring was observed (150°). At this point the remainder of the copper was added portionwise while keeping the temperature at 150°. After the addition was complete, the mixture was held at this temperature for an additional fifteen minutes.

At the end of this time the mixture was filtered hot with suction and the collected solids were washed thoroughly with acetone. The combined washings and filtrate were evaporated



CONFIDENTIAL  
NOLTR 62-175

in vacuo to remove the solvent and the solids remaining were taken up in acetone and treated with Norit to yield a dark red solution. The solution was evaporated to dryness and the residue titrated with ether. On filtering the slurry there was obtained about four grams of a tan amorphous solid; m. 169-72°. This product could not be purified by recrystallization.

The tan product was therefore hydrolyzed directly by heating it with 30 ml of concentrated sulfuric acid. At about 120° gas evolution (carbon dioxide) commenced. The mixture was heated at 120-40° until gas evolution ceased and then poured onto ice after cooling to about 50°. The yellow-brown product which separated was recrystallized from ethanol (Norit) to yield a yellow crystalline solid; m. 188-89.5°. No depression on admixture with an authentic sample of picramide.

CONFIDENTIAL  
NOLTR 62-175

REFERENCES

1. J. C. Dacons, H. G. Adolph, and M. J. Kamlet, NOLTR 62-82, Heat Resistant Explosives. XIII. Preparation and Properties of 3,3'-Diamino-2,2',4,4',6,6'-hexanitrobiphenyl (DIPAM), 16 May 1962.
2. W. W. Moyer, L. H. Bock, and R. Adams, J. Am. Chem. Soc., 52, 2054 (1930).
3. H. G. Adolph, J. C. Dacons, and M. J. Kamlet, NOLTR 62-32, Heat Resistant Explosives. XI. An Unusual Oxidation Reaction Leading to 3,3'-Dihydroxy-2,2',4,4',6,6'-hexanitrobiphenyl (Dipicric Acid, DIPA), 26 Feb 1962.
4. M. M. de Monchy, Rec. trav. chim., 53, 141 (1934).
5. See for example: P. E. Fanta, Chem. Reviews, 38, 139 (1946).
6. O. Diels and F. Bunzl, Ber., 38, 1486 (1905).
7. H. Lempert and R. Robinson, J. Chem. Soc., 1420 (1934).
8. J. C. Dacons and M. J. Kamlet, NavOrd Report (NOL) 6208, Heat Resistant Explosives. III. 1,3-Diamino-2,4,6-trinitrobenzene from 1,3-Dimethoxybenzene, 15 Dec 1958.
9. W. J. Hickinbottom, Reactions of Organic Compounds, 2nd edition, Longmans Green and Co., New York, page 376 (1950).
10. J. J. Blanksma, Rec. trav. chim., 23, 121 (1904).

CONFIDENTIAL  
NOLTR 62-175

DISTRIBUTION

	No. of Copies
Chief, Bureau of Naval Weapons, Department of the Navy, Washington 25, D. C.	
Attn: D-1 .....	1 1
DLI-3 .....	2
RRRE-5 .....	1
RUME-11 .....	1
RUME-323 .....	1
RMMP-21 .....	1
Chief, Office of Naval Research, Washington 25, D. C.	
Attn: Power Branch (Code 429) .....	1
Dr. Ralph Roberts .....	1
Chief of Ordnance, Department of the Army, Wash. 25, D.C.	
Attn: Technical Library .....	1
ODTM .....	1
Chief of Engineers, Department of the Army, Wash. 25, D.C.	
Attn: Research and Development Division .....	1
Commanding Officer, Naval Propellant Plant, Indian Head, Maryland	
Attn: Technical Library .....	1
Chemistry Research Section .....	1
Commanding Officer, Engineer Research and Development Laboratories, Fort Belvoir, Virginia	
Attn: Technical Library .....	1
Commanding Officer, U. S. Army Ordnance Missile Command, Redstone Arsenal, Huntsville, Alabama	
Attn: Technical Library .....	1
Rohm and Haas Division (SPIA C-36) .....	1
Commanding Officer, U. S. Naval Ordnance Laboratory, Corona, California	
Attn: Technical Library .....	1
Commanding Officer, Naval Weapons Station, Yorktown, Va.	
Attn: Research and Development Laboratory .....	1
Commanding General, Aberdeen Proving Ground, Maryland	
Attn: Ballistics Research Center .....	2
Commanding Officer, Department of the Army, Pantex Ordnance Plant, Amarillo, Texas .....	1
Commanding Officer, Lake City Arsenal, Ordnance Corps Department of the Army, Independence, Missouri	
Attn: Technical Library .....	1
Commander, U. S. Naval Weapons Laboratory, Dahlgren, Va.	
Attn: Technical Library .....	1
Commanding Officer, Holston Ordnance Works, U. S. Army, Kingsport, Tennessee	
Attn: Dr. Robert Robbins .....	1

CONFIDENTIAL

~~CONFIDENTIAL~~  
NOLTR 62-175

DISTRIBUTION  
(Cont'd)

	<u>No. of Copies</u>
Commander, Naval Ordnance Test Station, China Lake, California	
Attn: Chemistry Research Section .....	1
Dr. Martin H. Kaufman (Code 4544) .....	1
Headquarters, Air Proving Ground Center, USAF, ARDC, Eglin Air Force Base, Florida	
Attn: PGTRI, Technical Library .....	1
Headquarters, 6593rd Test Group (Dev.), AFSC, USAF Edwards Air Force Base, California	
Attn: (DGS) .....	1
Commanding Officer, Picatinny Arsenal, Dover, N. J.	
Attn: Library .....	2
Commanding General, U.S. Army Ordnance Arsenal, Frankford, Philadelphia 37, Penna.	
Attn: Propellant and Explosives Section .....	1
Director, Naval Research Laboratory, Washington 25, D.C.	
Attn: Chemistry Division .....	1
Director, Advanced Research Projects Agency, Pentagon, Rm. 3D159, Washington 25, D. C.	
Attn: Advanced Propellant Chemistry Office .....	1
Director, Special Projects Office, Department of the Navy, Washington 25, D. C.	
Attn: Technical Library .....	1
Solid Propellant Information Agency, Applied Physics Laboratory, Johns Hopkins Univ., Silver Spring, Md.	
Attn: Technical Library .....	3
Armed Forces Technical Information Agency, Arlington Hall Station, Arlington 12, Virginia	
Attn: TIPDR .....	10
Los Alamos Scientific Laboratory, P. O. Box 1663, Los Alamos, New Mexico (SPIA C-103)	
Attn: Reports Librarian for Dr. L. C. Smith .....	2
U. S. Naval Ammunition Depot, Navy No. Six Six (66), c/o FPO, San Francisco, California	
Attn: Quality Evaluation Laboratory .....	2
University of California, Lawrence Radiation Laboratory, P.O. Box 808, Livermore, California	
Attn: Dr. Gus Dorough, Chemistry Division .....	1
Aerojet-General Corporation, Azusa, California	
Attn: Dr. J. R. Fischer (SPIA C-1) .....	1
Aerojet-General Corporation, P.O. Box 1947, Sacramento, California (SPIA C-72)	
Attn: Dr. Marvin H. Gold .....	1

~~CONFIDENTIAL~~

~~CONFIDENTIAL~~  
NOLTR 62-175

DISTRIBUTION  
(Cont'd)

No. of  
Copies

American Cyanamid Company, P.O. Box 270, New Castle, Pa. Attn: L. V. Clark, Director (Contr. NOW-60-0745c) Explosives Research and Development .....	1
Minnesota Mining and Manufacturing Company 2301 Hudson Rd., St. Paul 19, Minn. (ARPA Proj. 201-2) Attn: Dr. M. H. Friedman, Central Research Dep't.)...	1
Minnesota Mining and Manufacturing Co. 900 Bush Avenue St. Paul 6, Minnesota (SPIA C-94) Attn: Dr. Wm. S. Friedlander Via: H. C. Zeman, Security Administrator .....	1
Hercules Powder Co., Explosives Dep't., Delaware Trust Bldg., Wilmington 99, Delaware Attn: A. M. Ball (SPIA C-18) .....	1
Purdue University, Dep't. of Chemistry Lafayette, Indiana (SPIA C-33) Attn: Dr. Henry Feuer .....	1
Esso Research and Engineering Company, Chemicals Research Division, P.O. Box 51, Linden, N. J. (SPIA C-95) Attn: Dr. J. P. Longwell Via: Chief, N.Y. Ordnance District .....	1
Lockheed Aircraft Corporation, Missiles and Space Div. Attn: Technical Library (Contract Nord 17017) Via: Bur. of Naval Weapons Rep. (Code SpL 31) Special Proj. Office, P.O. Box 504, Sunnyvale, California .....	1

~~CONFIDENTIAL~~

CONFIDENTIAL

CATALOGING INFORMATION FOR LIBRARY USE

BIBLIOGRAPHIC INFORMATION					
	DESCRIPTORS	CODES		DESCRIPTORS	CODES
SOURCE	NOL technical report	NOLTR	SECURITY CLASSIFICATION AND CODE COUNT	Confidential - 21	0021
REPORT NUMBER	62-175	620175	CIRCULATION LIMITATION		
REPORT DATE	24 September 1962	0962	CIRCULATION LIMITATION OR BIBLIOGRAPHIC		
			BIBLIOGRAPHIC (SUPPL., VOL., ETC.)		

SUBJECT ANALYSIS OF REPORT

	DESCRIPTORS	CODES	DESCRIPTORS	CODES	DESCRIPTORS	CODES
Heat-resistant	HEAR		Derivative	DEBI		
Explosives	EXPL		Ammonia	AMMO		
Diamino	DIAM		Alcohol	ALCO		
Hexa	HEXA		Aniline	ANIL		
Nitro	NIRO		Methyl	METY		
Diphenyl	DIPH		Phenyl	PHEN		
Synthesis	SYNS		Urethane	URET		
Bromo	BROO		Bis	BISE		
Anisole	ANSL		Picramide	FORM		
Trinitro	TRIT					
Dimethoxy	DIMH					
Conversion	CNVE					

CONFIDENTIAL

<p>Naval Ordnance Laboratory, White Oak, Md. (NOL technical report 62-175) HEAT RESISTANT EXPLOSIVES. XV. (U) ALTERNATE SYNTHETIC ROUTES TO 3,3'-DIAMINO-2,2',4,4',6,6'-HEXANTROBIPHENYL, DIPAM (C), by Robert E. Oesterling and others. 24 Sept. 1962. 11p. diagr. NOL task FR-44 CONFIDENTIAL</p> <p>DIPAM has been prepared from m-bromoanisole by a new three step synthesis involving the following reactions: m-bromoanisole to trinitro-m-bromoanisole which was in turn converted to dimethoxy hexanitrobiphenyl via the Ullmann reaction. Conversion of the dimethoxy derivative to DIPAM was readily achieved by treating it with ammonia in alcohol.</p> <p>Abstract card is confidential</p> <p>DOWNGRADED AT 3 YEAR INTERVALS: DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10.</p>	<p>1. Explosives, Heat resistant 3,3'-Diamino-2,2',4,4',6,6'-hexanitrobiphenyl</p> <p>2. Explosives-Production Title: Title: Alternate synthetic routes to DIPAM</p> <p>III. Oesterling Robert E. IV. Project</p>
<p>Naval Ordnance Laboratory, White Oak, Md. (NOL technical report 62-175) HEAT RESISTANT EXPLOSIVES. XV. (U) ALTERNATE SYNTHETIC ROUTES TO 3,3'-DIAMINO-2,2',4,4',6,6'-HEXANTROBIPHENYL, DIPAM (C), by Robert E. Oesterling and others. 24 Sept. 1962. 11p. diagr. NOL task FR-44 CONFIDENTIAL</p> <p>DIPAM has been prepared from m-bromoanisole by a new three step synthesis involving the following reactions: m-bromoanisole to trinitro-m-bromoanisole which was in turn converted to dimethoxy hexanitrobiphenyl via the Ullmann reaction. Conversion of the dimethoxy derivative to DIPAM was readily achieved by treating it with ammonia in alcohol.</p> <p>Abstract card is confidential</p> <p>DOWNGRADED AT 3 YEAR INTERVALS: DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10.</p>	<p>1. Explosives, Heat resistant 3,3'-Diamino-2,2',4,4',6,6'-hexanitrobiphenyl</p> <p>2. Explosives-Production Title: Title: Alternate synthetic routes to DIPAM</p> <p>III. Oesterling Robert E. IV. Project</p>
<p>Naval Ordnance Laboratory, White Oak, Md. (NOL technical report 62-175) HEAT RESISTANT EXPLOSIVES. XV. (U) ALTERNATE SYNTHETIC ROUTES TO 3,3'-DIAMINO-2,2',4,4',6,6'-HEXANTROBIPHENYL, DIPAM (C), by Robert E. Oesterling and others. 24 Sept. 1962. 11p. diagr. NOL task FR-44 CONFIDENTIAL</p> <p>DIPAM has been prepared from m-bromoanisole by a new three step synthesis involving the following reactions: m-bromoanisole to trinitro-m-bromoanisole which was in turn converted to dimethoxy hexanitrobiphenyl via the Ullmann reaction. Conversion of the dimethoxy derivative to DIPAM was readily achieved by treating it with ammonia in alcohol.</p> <p>Abstract card is confidential</p> <p>DOWNGRADED AT 3 YEAR INTERVALS: DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10.</p>	<p>1. Explosives, Heat resistant 3,3'-Diamino-2,2',4,4',6,6'-hexanitrobiphenyl</p> <p>2. Explosives-Production Title: Title: Alternate synthetic routes to DIPAM</p> <p>III. Oesterling Robert E. IV. Project</p>
<p>Naval Ordnance Laboratory, White Oak, Md. (NOL technical report 62-175) HEAT RESISTANT EXPLOSIVES. XV. (U) ALTERNATE SYNTHETIC ROUTES TO 3,3'-DIAMINO-2,2',4,4',6,6'-HEXANTROBIPHENYL, DIPAM (C), by Robert E. Oesterling and others. 24 Sept. 1962. 11p. diagr. NOL task FR-44 CONFIDENTIAL</p> <p>DIPAM has been prepared from m-bromoanisole by a new three step synthesis involving the following reactions: m-bromoanisole to trinitro-m-bromoanisole which was in turn converted to dimethoxy hexanitrobiphenyl via the Ullmann reaction. Conversion of the dimethoxy derivative to DIPAM was readily achieved by treating it with ammonia in alcohol.</p> <p>Abstract card is confidential</p> <p>DOWNGRADED AT 3 YEAR INTERVALS: DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10.</p>	<p>1. Explosives, Heat resistant 3,3'-Diamino-2,2',4,4',6,6'-hexanitrobiphenyl</p> <p>2. Explosives-Production Title: Title: Alternate synthetic routes to DIPAM</p> <p>III. Oesterling Robert E. IV. Project</p>